



Introduction

Foot and ankle surgery is a specialised area of surgery. The surgery can be quite challenging and every patient is different with unique needs and goals.

You will need to follow instructions, perform appropriate exercises and modify your activities during your healing process.

This requires patience, persistence and a desire to get better. If you are unable to complete the post-operative instructions it will affect your results and you should consider alternative treatments.

Whenever surgery is considered we always try to minimise any risks. You are already on this path by consulting an experienced specialist who exclusively treats foot and ankle disorders. The body is a very complex and varied structure so although we aim for perfection no specialist can give perfect results every time.

Successful results require a contribution from you

This information package is intended to provide you with pre-operative, surgical and post-operative instructions in regard to your surgical procedure.

It is important that you read all of this information carefully and bring it with you when you come to hospital. If you have any questions or are not certain about the benefits, risks and limitations of your treatment ask your surgeon.

What do I take to Hospital?

- Your X-rays.
- Documentation about your Health Fund.
- Pathology Results.
- Medication.
- Any letters from doctors.
- This information pack.
- Any questions that you have written down and wish to ask.

Will I need someone to take me home?

You must make arrangements for a responsible adult to take you home after your surgery. This is advisable if you have either had a day procedure or if you are in hospital for a period of time. It is strongly suggested that you have someone stay with you during the first 24 hours or until you are able to get around safely.

What should I wear?

Wear loose fitting clothes that are easy to put on and will fit over bulky bandages or surgical dressings. Leave any jewellery and valuables at home. Please remove make up and nail polish. Shower either the night before or the morning of your surgery. Pay particular care that your feet and nails are clean.

Local Anaesthetic Block for Pain Management

A local anaesthetic block may be given to you during surgery. This will produce numbness around the nerves in the region of your surgical procedure. It will provide you with pain relief for approximately 8–12 hours post-operatively enabling you to be comfortable and allow you to sleep after your surgery.

Discharge Home

- Plan to **rest** after surgery to minimise bleeding and swelling. Your particular instruction sheet will give you a specific period of time.
- Have someone who is responsible available to take you home, as **you will not be allowed to drive**.
- Once you arrive home wait until you are hungry before eating. Begin with a light meal such as a sandwich/fruit/tea/coffee or juice. A heavy meal can cause nausea and vomiting after an operation.
- Take your medication for pain as directed.
- A physiotherapist may see you if required while you are in hospital to give you instruction as to walking, weight bearing, exercises and use of any aids. (eg. Crutches).
- Once you can demonstrate that you can safely use crutches, able to negotiate stairs and pain relief is adequate, then you can go home.

Post-Operative Instructions

These instructions are important in helping you rehabilitate from your surgery. Please follow them carefully. If there is anything you do not understand, please ask.

Rest

It is important that you rest and keep your foot elevated as much as possible for the first 7–10 days to encourage healing and discourage wound break down.

Patient Surgical Information

Foot and Ankle Surgery



Mercy Private Hospital

Post-Operative Instructions

Pain Management

Pain is better treated before it arrives. A regular regime of pain killers taken in the post-operative phase is very important. It's recommended that you take regular analgesic, Digesic or Panadeine Forte 4-6 hourly initially, whilst pain is moderate to severe, however after that period it is recommended that you take regular Panadol up until your post-operative visit.

Please take analgesic 2 hours prior to your post-operative appointment.

Two hours before going to bed take two analgesic to help you sleep.

If you have any problems taking any of these medications please let the medical staff know.

There is no need for you to be in constant pain.

Elevation

Swelling can delay wound healing and cause the wound to break down. Swelling can be due to over activity and not keeping your foot elevated and this can also increase the risk of infection. Place your operated leg on two pillows or a bean bag so that it is above the level of your hip when you are lying down. The use of ice packs (or frozen peas, beans etc.) can be a very effective way of reducing swelling. Place the ice pack in a towel over the ankle or foot for 20 minutes 3-4 times per day.

Exercise

While you are resting it is important that you do passive gentle movement exercises. This is to encourage your blood flow, muscle strengthening and prevention of joint stiffness. The exercise information given to you by your surgeon or the physiotherapist is designed to help you with your rehabilitation and to gain the best results after surgery. Continue with your exercises once you are home and up until your post-operative appointment.

Dressings

- Keep your bandages/cast clean and dry. Leave in tact until your post-operative appointment.
- When showering/bathing, wrap the dressings or cast in a plastic kitchen or garbage bag and seal the top of the bag above your dressings with tape.

Post-Operative Problems

If after reading and following the post-operative instructions you are experiencing any of the following problems please call the rooms:

Mr Blackney's Rooms (03) 9417 0762

Mr Chin's Rooms (03) 9928 6450

Infection

- Wounds feel hot and tender.
- Unusual discharge from your wound or dressings or an odour from your dressings.
- Fever may be present and a general feeling of unwell.

Swelling

- Swelling is not reduced with elevation of the leg.
- A change in colour of your toes or coldness.
- The calf muscle in your lower leg of the treated foot swells or is painful.

Bleeding

- Your dressings are showing excessive amount of fresh blood.
- Elevation and application of pressure dressings doesn't stop the bleeding.

Pain

- The medication that has been prescribed for you is not helping control the pain while you have been resting with your limb elevated.

Medication

- You are experiencing ill effects due to the medication that has been prescribed for pain eg. Nausea, vomiting, gastric discomfort.

Mark C. Blackney

MBBS BSc FRACS
Orthopaedic Surgeon

Phone (03) 9417 0762

Fax (03) 9417 0433

Terence Chin

MBBS FRACS
Orthopaedic Surgeon

Phone (03) 9928 6450

Fax (03) 9417 0433

The Park Clinic
Specialised Orthopaedics

www.theparkclinic.com.au